

**PATENT APPLICATION TRANSMITTAL LETTER**

(Large Entity)

Docket No.

1038-971 MIS:jb

TO THE ASSISTANT COMMISSIONER FOR PATENTS

Transmitted herewith for filing under 35 U.S.C. 111 and 37 C.F.R. 1.53 is the patent application of:

Andrew D. Murdin; and Pamela L. Dunn

**IMMUNOGENIC COMPOSITIONS FOR PROTECTION AGAINST CHLAMYDIAL INFECTION**

Enclosed are:

- ☐ Certificate of Mailing with Express Mail Mailing Label No.
- ☒ **Ten (10)** sheets of drawings.
- ☐ A certified copy of a application.
- ☒ Declaration ☐ Signed. ☒ Unsigned.
- ☐ Power of Attorney
- ☐ Information Disclosure Statement
- ☐ Preliminary Amendment
- ☐ Other:

**CLAIMS AS FILED**

For	#Filed	#Allowed	#Extra	Rate	Fee
<b>Total Claims</b>	23	- 20 =	3	x \$18.00	\$54.00
<b>Indep. Claims</b>	2	- 3 =	0	x \$78.00	\$0.00
<b>Multiple Dependent Claims (check if applicable)</b> <input type="checkbox"/>					\$0.00
<b>BASIC FEE</b>					\$760.00
<b>TOTAL FILING FEE</b>					\$814.00

- ☒ A check in the amount of **\$814.00** to cover the filing fee is enclosed.
- ☒ The Commissioner is hereby authorized to charge and credit Deposit Account No. as described below. A duplicate copy of this sheet is enclosed.
- ☐ Charge the amount of as filing fee.
- ☒ Credit any overpayment.
- ☒ Charge any additional filing fees required under 37 C.F.R. 1.16 and 1.17.
- ☐ Charge the issue fee set in 37 C.F.R. 1.18 at the mailing of the Notice of Allowance, pursuant to 37 C.F.R. 1.311(b).

Dated: September 2, 1999

*Michael I. Stewart*  
Signature  
Michael I. Stewart (24,973)

cc:

TITLE OF INVENTION  
**IMMUNOGENIC COMPOSITIONS FOR PROTECTION  
AGAINST CHLAMYDIAL INFECTION**

FIELD OF THE INVENTION

5           The present invention relates to immunogenic compositions for protection against disease caused by *Chlamydia* infection in mammals, including humans.

BACKGROUND OF THE INVENTION

10           Chlamydiae are procaryotes. They exhibit morphologic and structural similarities to gram negative bacteria, including a trilaminar outer membrane, which contains lipopolysaccharide and several membrane proteins. Chlamydiae are differentiated from other bacteria by their morphology and by a unique developmental cycle. They are obligate intracellular parasites with a unique biphasic life cycle consisting of a  
15           metabolically inactive but infectious extracellular stage and a replicating but non-infectious intracellular stage. The replicative stage of the life-cycle takes place within a membrane-bound inclusion which sequesters the bacteria away from the cytoplasm of the infected host cell.

20           Because chlamydiae are small and multiply only within susceptible cells, they were long thought to be viruses. However, they have many characteristics in common with other bacteria: (1) they contain both DNA and RNA, (2) they divide by binary fission, (3) their cell envelopes resemble those of other gram-negative bacteria, (4) they contain ribosomes similar to those of other bacteria, and (5) they are susceptible  
25           to various antibiotics. Chlamydiae can be seen in the light microscope, and the genome is about one-third the size of the *Escherichia coli* genome.

30           Many different strains of chlamydiae have been isolated from birds, man and other mammals, and these strains can be distinguished on the basis of host range, virulence, pathogenesis, and antigenic composition.

09391606 1090799

There is strong homology of DNA within each species, but surprisingly little between species, suggesting long-standing evolutionary separation.

*C. trachomatis* has a high degree of host specificity, being almost completely limited to man, and causes ocular and genitourinary infections of widely varying severity. In contrast, *C. psittaci* strains are rare in man but are found in a wide range of birds and also in wild, domestic, and laboratory mammals, where they multiply in cells of many organs.

*C. pneumoniae* is a common human pathogen, originally described as the TWAR strain of *C. psittaci*, but subsequently recognized to be a new species. *C. pneumoniae* is antigenically, genetically, and morphologically distinct from other *Chlamydia* species (*C. trachomatis*, *C. pecorum* and *C. psittaci*). It shows 10% or less DNA sequence homology with either of *C. trachomatis* or *C. psittaci* and so far appears to consist of only a single strain, TWAR.

*C. pneumoniae* is a common cause of community acquired pneumonia, less frequent only than *Streptococcus pneumoniae* and *Mycoplasma pneumoniae* (refs. 1 and 2 - Throughout this application, various references are referred to in parenthesis to more fully describe the state of the art to which this invention pertains. Full bibliographic information for each citation is found at the end of the specification, immediately preceding the claims. The disclosure of these references are hereby incorporated by reference into the present disclosure). *C. pneumoniae* can also cause upper respiratory tract symptoms and disease, including bronchitis and sinusitis (refs. 1 to 4). The great majority of the adult population (over 60%) has antibodies to *C. pneumoniae* (ref. 5), indicating past infection which was unrecognized or asymptomatic.

*C. pneumoniae* infection usually presents as an acute respiratory disease (*i.e.*, cough, sore throat, hoarseness, and fever; abnormal chest sounds on auscultation). For most patients, the cough persists for 2 to 6 weeks, and recovery is slow. In approximately 10% of these cases, upper respiratory tract infection is followed by bronchitis or pneumonia. Furthermore, during a *C. pneumoniae* epidemic, subsequent co-infection

with pneumococcus has been noted in about half of these pneumonia patients, particularly in the infirm and the elderly. As noted above, there is more and more evidence that *C. pneumoniae* infection is also linked to diseases other than respiratory infections.

5       The reservoir for the organism is presumably people. In contrast to *C. psittaci* infections, there is no known bird or animal reservoir. Transmission has not been clearly defined, but may result from direct contact with secretions, from fomites, or from airborne spread. There is a long incubation period, which may last for many months. Based on  
10 analysis of epidemics, *C. pneumoniae* appears to spread slowly through a population (case-to-case interval averaging 30 days) because infected persons are inefficient transmitters of the organism. Susceptibility to *C. pneumoniae* is universal. Reinfections occur during adulthood, following the primary infection as a child. *C. pneumoniae* appears to be an  
15 endemic disease throughout the world, noteworthy for superimposed intervals of increased incidence (epidemics) that persist for 2 to 3 years. *C. trachomatis* infection does not confer cross-immunity to *C. pneumoniae*. Infections are easily treated with oral antibiotics, tetracycline or erythromycin (2 g/d, for at least 10 to 14 d). A recently developed drug,  
20 azithromycin, is highly effective as a single-dose therapy against chlamydial infections.

In most instances, *C. pneumoniae* infection is mild and without complications, and up to 90% of infections are subacute or unrecognized. Among children in industrialized countries, infections have been thought to  
25 be rare up to the age of 5 years, although a recent study has reported that many children in this age group show PCR evidence of infection despite being seronegative, and estimates a prevalence of 17 to 19% in 2 to 4 years old (ref. 6). In developing countries, the seroprevalence of *C. pneumoniae* antibodies among young children is elevated, and there  
30 are suspicions that *C. pneumoniae* may be an important cause of acute lower respiratory tract disease and mortality for infants and children in tropical regions of the world.

09391606 "090799

From seroprevalence studies and studies of local epidemics, the initial *C. pneumoniae* infection usually happens between the ages of 5 and 20 years. In the USA, for example, there are estimated to be 30,000 cases of childhood pneumonia each year caused by *C. pneumoniae*. Infections may cluster among groups of children or young adults (e.g., school pupils or military conscripts).

*C. pneumoniae* causes 10 to 25% of community-acquired lower respiratory tract infections (as reported from Sweden, Italy, Finland, and the USA). During an epidemic, *C. pneumoniae* infection may account for 50 to 60% of the cases of pneumonia. During these periods, also, more episodes of mixed infections with *S. pneumoniae* have been reported.

Reinfection during adulthood is common; the clinical presentation tends to be milder. Based on population seroprevalence studies, there tends to be increased exposure with age, which is particularly evident among men. Some investigators have speculated that a persistent, asymptomatic *C. pneumoniae* infection state is common.

In adults of middle age or older, *C. pneumoniae* infection may progress to chronic bronchitis and sinusitis. A study in the USA revealed that the incidence of pneumonia caused by *C. pneumoniae* in persons younger than 60 years is 1 case per 1,000 persons per year; but in the elderly, the disease incidence rose three-fold. *C. pneumoniae* infection rarely leads to hospitalization, except in patients with an underlying illness.

Of considerable importance is the association of atherosclerosis and *C. pneumoniae* infection. There are several epidemiological studies showing a correlation of previous infections with *C. pneumoniae* and heart attacks, coronary artery and carotid artery disease (refs. 7 to 11). Moreover, the organisms has been detected in atheromas and fatty streaks of the coronary, carotid, peripheral arteries and aorta (refs. 12 to 16). Viable *C. pneumoniae* has been recovered from the coronary and carotid artery. (refs, 17, 18). Furthermore, it has been shown that *C. pneumoniae* can induce changes of atherosclerosis in a rabbit model (ref. 19). Taken together, these results indicate that it is highly probable

that *C. pneumoniae* can cause atherosclerosis in humans, though the epidemiological importance of chlamydial atherosclerosis remains to be demonstrated.

5 A number of recent studies have also indicated an association between *C. pneumoniae* infection and asthma. Infection has been linked to wheezing, asthmatic bronchitis, adult-onset asthma and acute exacerbation of asthma in adults, and small-scale studies have shown that prolonged antibiotic treatment was effective at greatly reducing the severity of the disease in some individuals (refs. 20 to 25).

10 In light of these results, a protective vaccine against disease caused by *C. pneumoniae* infection would be of considerable importance. There is not yet an effective vaccine for human *C. pneumoniae* infection. Nevertheless, studies with *C. trachomatis* and *C. psittaci* indicate that this is an attainable goal. For example, mice which have recovered from a  
15 lung infection with *C. trachomatis* are protected from infertility induced by a subsequent vaginal challenge (ref. 26). Similarly, sheep immunized with inactivated *C. psittaci* were protected from subsequent chlamydial-induced abortions and stillbirths (ref. 27). Protection from chlamydial infections has been associated with Th1 immune responses, particularly the induction of  
20 INF $\gamma$ -producing CD4<sup>+</sup> T cells (ref. 28). The adoptive transfer of CD4<sup>+</sup> cell lines or clones to nude or SCID mice conferred protection from challenge or cleared chronic disease (refs. 29, 30) and *in vivo* depletion of CD4<sup>+</sup> T cells exacerbated disease post-challenge (refs. 31, 32). However, the presence of sufficiently high titres of neutralizing antibody at mucosal  
25 surfaces can also exert a protective effect (ref. 33).

The extent of antigenic variation within the species *C. pneumoniae* is not well characterized. Serovars of *C. trachomatis* are defined on the basis of antigenic variation in major outer membrane proteins (MOMP), but published *C. pneumoniae* MOMP gene sequences show no variation  
30 between several diverse isolates of the organism (refs. 34, 35, 36). Regions of the protein known to be conserved in other chlamydial MOMP are conserved in *C. pneumoniae* (refs. 34, 35). One study has described a

strain of *C. pneumoniae* with a MOMP of greater than usual molecular weight, but the gene for this has not been sequenced (ref. 1). Partial sequences of outer membrane protein 2 from nine diverse isolates were also found to be invariant (ref. 17). The genes for HSP60 and HSP70 show little variation from other chlamydial species, as would be expected. The gene encoding a 76 kDa antigen has been cloned from a single strain of *C. pneumoniae*. It has no significant similarity with other known chlamydial genes (ref. 4).

Many antigens recognized by immune sera to *C. pneumoniae* are conserved across all chlamydiae, but 98kDa, 76 kDa and 54 kDa proteins may be *C. pneumoniae*-specific (refs. 2, 4, 37). Immunoblotting of isolates with sera from patients does show variation of blotting patterns between isolates, indicating that serotypes *C. pneumoniae* may exist (refs. 1, 17). However, the results are potentially confounded by the infection status of the patients, since immunoblot profiles of a patient's sera change with time post-infection. An assessment of the number and relative frequency of any serotypes, and the defining antigens, is not yet possible.

Thus, a need remains for effective compositions for preventing and treating *Chlamydia* infections.

#### SUMMARY OF THE INVENTION

The present invention provides a novel approach to immunizing against Chlamydial infection based on nucleic acid immunization. It has surprisingly been found that the administration of a combination of nucleotide sequences encoding two different chlamydial proteins provides an enhanced protection efficacy.

Accordingly, in one aspect of the present invention, there is provided an immunogenic composition for *in vivo* administration to a host for the generation in the host of a protective immune response against Chlamydial infection, comprising a first vector comprising a first nucleotide sequence encoding a major outer membrane protein (MOMP) of a strain of *Chlamydia* and a first promoter sequence operatively coupled to said first nucleotide sequence for expression of said MOMP in the host; a second

vector comprising a second nucleotide sequence encoding a 76 kDa protein of a strain of *Chlamydia* and a second promoter sequence operatively coupled to said second nucleotide sequence for expression of said 76 kDa protein in the host; and a pharmaceutically-acceptable carrier therefor.

The first nucleotide sequence may encode a MOMP from any strain of *Chlamydia*, preferably from *C. pneumoniae* but also including *C. trachomatis*. The second nucleotide sequence encoding the MOMP protein of *C. pneumoniae* may have SEQ ID No: 12, 13 or 14 or may encode a MOMP having a SEQ ID No: 15 or 16.

The first promoter which is employed may be a cytomegalovirus promoter, although any other convenient promoter may be employed.

The second nucleotide sequence may encode a 76 kDa protein from any strain of *Chlamydia*, preferably from *C. pneumoniae* but also including *C. trachomatis*. The second nucleotide sequence encoding the 76 kDa protein of *C. pneumoniae* may have SEQ ID No: 1, 2, 3 or 4. The second nucleotide sequence may encode a 76 kDa protein having a molecular size of about 35 kDa and having SEQ ID No: 7 or may encode a 76 kDa protein having a molecular size of about 60 kDa and having SEQ ID No: 8 or 9.

The second promoter which is employed may be a cytomegalovirus promoter, although any other convenient promoter may be employed.

The first vector preferably comprises a plasmid vector and specifically may be pCAMOMP. Similarly the second vector preferably comprises a plasmid vector and specifically may be pCA76kDa. Most preferably, both the first and second vectors are plasmid vectors and specifically the combination of pCAMOMP and pCA76kDa.

The two vectors are used in an immunogenic composition along with any convenient pharmaceutically-acceptable carrier. As noted above, the uses of the combination of two vectors produces an enhanced protection efficacy in comparison to the individual vectors alone. Accordingly, the first and second vectors preferably are present in the



The present invention, in a further aspect thereof, provides a method of immunizing a host against disease caused by infection with a strain of *Chlamydia*, which comprises administering to the host, which may be a human host, an effective amount of an immunogenic composition provided herein. The immunogenic composition preferably is administered intranasally.

## 10

Figure 1 shows the nucleotide sequence of *C. pneumoniae* 76kDa gene (SEQ ID No: 1 - complete sequence; SEQ ID No: 2 - 5' encoding region; SEQ ID No: 3 - 3' encoding region including Myc and His encoding regions; SEQ ID No: 4 - 3' encoding region excluding Myc and His encoding regions; SEQ ID No: 5 - Myc encoding region; SEQ ID No: 6 - His encoding region) and the deduced amino acid sequences of two open reading frames of the 76kDa protein (SEQ ID NO: 7 - upstream reading frame; SEQ ID No: 8 - downstream reading frame including Myc and His regions; SEQ ID No: 9 - downstream reading frame excluding Myc and His regions; SEQ ID No: 10 - Myc region; SEQ ID No: 11 - His region):

Figure 3 shows the nucleotide sequence of the *C. pneumoniae* MOMP gene (SEQ ID No: 12 - complete sequence; SEQ ID No: 13 - encoding sequence including Myc and His encoding regions; SEQ ID No: 14 - encoding sequence excluding Myc and His encoding regions) and the deduced amino acid sequence of the MOMP protein (SEQ ID No: 15 - including Myc and His regions; SEQ ID No: 16 - excluding Myc and His regions);

Figure 4 shows a scheme of the construction of plasmid pCAMOMP; and

Figure 5 illustrates the protective efficacy against *C. pneumoniae* lung challenge in Balb/c mice following DNA immunization with pCAMOMP plus pCA76kDa, in comparison to controls, wherein the individual data points (open diamonds) are shown for each animal, as well as the mean (solid squares) and standard deviation for each group.

#### GENERAL DESCRIPTION OF INVENTION

As noted above, the present invention is directed to protecting a host against chlamydial infection by administering to the host an immunogenic composition containing two vectors, preferably plasmid vectors, each of which contains nucleotide sequence encoding a different protein of a strain of *Chlamydia*.

To illustrate the invention, a first plasmid vector was constructed containing the MOMP gene from *C. pneumoniae* and a second plasmid vector was constructed containing the 76 kDa protein gene from *C. pneumoniae*. While the invention is illustrated by the use of such plasmid vectors, other vectors containing such genes may be employed for administration to the host for expression of the encoded proteins in the host. Such other vectors may include live viral vectors, such as adenoviruses, alphaviruses including Semliki Forest virus and poxviruses including avipox and canary pox viruses as well as bacterial vectors, such as *Shigella*, *Salmonella*, *Vibrio cholerae*, *Lactobacillus*, Bacille Bilié de Calmette-Guérin (BCG) and *Streptococcus*.

One of the vectors employed herein contains a nucleic acid molecule which codes for a Chlamydial protein known in the art as the "76 kDa protein" (ref. 4). The latter terminology is utilized herein to refer to the protein identified in the art. Research has determined that the encoding nucleotide sequence for this protein in fact encodes two opening reading frames, one encoding a protein of approximately 35 kDa in length (SEQ ID No: 7) and the other encoding a protein of approximately 60 kDa in length (SEQ ID No: 9).

It has been found that, if the complete nucleotide sequence (SEQ ID No: 1) is incorporated into a suitable expression vector, then only the 35

kDa protein is expressed. If, however, the nucleotide sequence encoding the 60 kDa protein alone (SEQ ID No: 4) is incorporated into a suitable expression vector, then that protein also can be expressed. Both proteins have been found to be immunogenic and protective with the 35 kDa protein exhibiting a stronger protective effect than the 60 kDa protein (United States Patent Application No. 60/132,270 filed May 3, 1999; United States Patent Application No. 60/141,276 filed June 30, 1999, assigned to the Assignee hereof and the disclosures of which are incorporated herein by reference).

Any convenient plasmid vector may be used for the MOMP gene and the 76 kDa protein gene, such as the pcDNA3.1 expression vector (Invitrogen, San Diego, CA, USA) containing the cytomegalovirus promoter. Schemes for construction of the pCA76kDa plasmid vector of 8594 bp size and of the pCAMOMP plasmid vector of 7.6 kb in size, which include downstream DNA sequences coding for Myc and His tags, are shown in Figures 2 and 4 respectively and described in detail below.

The respective plasmids are formulated into an immunogenic composition in conjunction with a suitable pharmaceutically-acceptable carrier for administration to a host, such as a human host. The immunogenic composition may be administered in any convenient manner to the host, such as intramuscularly or intranasally, although other routes of administration may be used, as discussed below. The data presented herein and described in detail below demonstrates that DNA immunization with both the *C. pneumoniae* MOMP and 76 kDa protein genes elicits a strong protective immune response. The effect which is obtained is achieved without the use of adjuvant or other stimulation of immune response, such as cardiotoxin, although such materials may be used, if desired, as discussed below. In addition, the use of immunomodulation is not excluded from the scope of the invention. For example, it may be desirable to coadminister DNA that expresses immunoregulator cytokines (ref. 38).

As may be seen from the data below, by utilizing both the MOMP gene and the 76 kDa protein gene, there is obtained a protective immune response which is significantly greater than that achieved using the individual genes alone. The coadministration of the two genes does not  
 5 result in any interference to the immune response of the individual genes.

There has previously been described in WO 98/02546, assigned to University of Manitoba and the disclosure of which is incorporated herein by reference, the use of the MOMP gene for DNA immunization. The improved results obtained herein using a combination of the MOMP gene  
 10 and the 76 kDa protein gene demonstrate the use of multiple antigen genes from chlamydiae to augment the level of protective immunity achieved by DNA immunization. These results are more encouraging than those obtained using recombinant MOMP protein or synthetic peptides as the immunogen.

Nucleotide sequences, *e.g.*, DNA molecules, can easily be retrieved by polymerase chain reaction (PCR) amplification of genomic bacterial DNA extracted by conventional methods. This involves the use of synthetic oligonucleotide primers matching upstream and downstream of the 5' and 3' ends of the encoding domain. Suitable primers can be  
 15 designed according to the nucleotide sequence information provided. Typically, a primer can consist of 10 to 40, preferably 15 to 25 nucleotides. It may be also advantageous to select primers containing C and G nucleotides in a proportion sufficient to ensure efficient hybridization; *e.g.*, an amount of C and G nucleotides of at least 40%, preferably 50% of the  
 20 total nucleotide amount.

It is clearly apparent to one skilled in the art that the various embodiments of the present invention have many applications in the fields of vaccination and treatment of chlamydial infection. A further non-limiting discussion of such uses is further presented below.

### 30 1. Vaccine Preparation and Use

Immunogenic compositions, suitable to be used as vaccines, may be prepared from the MOMP gene and the 76 kDa protein gene and vectors as

disclosed herein. The vaccine elicits an immune response in a subject which includes the production of anti-MOMP and anti-76 kDa protein antibodies. Immunogenic compositions, including vaccines, containing the nucleic acid may be prepared as injectables, in physiologically-acceptable liquid solutions  
 5 or emulsions for polynucleotide administration.

The nucleic acid may be associated with liposomes, such as lecithin liposomes or other liposomes known in the art, as a nucleic acid liposome (for example, as described in WO 93/24640) or the nucleic acid may be associated with an adjuvant, as described in more detail below. Liposomes  
 10 comprising cationic lipids interact spontaneously and rapidly with polyanions, such as DNA and RNA, resulting in liposome/nucleic acid complexes that capture up to 100% of the polynucleotide. In addition, the polycationic complexes fuse with cell membranes, resulting in an intracellular delivery of polynucleotide that bypasses the degradative enzymes of the lysosomal  
 15 compartment.

Published PCT application WO 94/27435 describes compositions for genetic immunization comprising cationic lipids and polynucleotides. Agents which assist in the cellular uptake of nucleic acid, such as calcium ions, viral proteins and other transfection facilitating agents, may advantageously be  
 20 used.

Polynucleotide immunogenic preparations may also be formulated as microcapsules, including biodegradable time-release particles. Thus, U.S. Patent 5,151,264 describes a particulate carrier of a phospholipid/glycolipid/polysaccharide nature that has been termed Bio  
 25 Vecteurs Supra Moléculaires (BVSM). The particulate carriers are intended to transport a variety of molecules having biological activity in one of the layers thereof.

U.S. Patent 5,075,109 describes encapsulation of the antigens trinitrophenylated keyhole limpet hemocyanin and staphylococcal enterotoxin  
 30 B in 50:50 poly (DL-lactide-co-glycolide). Other polymers for encapsulation are suggested, such as poly(glycolide), poly(DL-lactide-co-glycolide), copolyoxalates, polycaprolactone, poly(lactide-co-caprolactone),

poly(esteramides), polyorthoesters and poly(8-hydroxybutyric acid), and polyanhydrides.

Published PCT application WO 91/06282 describes a delivery vehicle comprising a plurality of bioadhesive microspheres and antigens. The  
5 microspheres being of starch, gelatin, dextran, collagen or albumin. This delivery vehicle is particularly intended for the uptake of vaccine across the nasal mucosa. The delivery vehicle may additionally contain an absorption enhancer.

The vectors may be mixed with pharmaceutically acceptable  
10 excipients which are compatible therewith. Such excipients may include, water, saline, dextrose, glycerol, ethanol, and combinations thereof. The immunogenic compositions and vaccines may further contain auxiliary substances, such as wetting or emulsifying agents, pH buffering agents, or adjuvants to enhance the effectiveness thereof. Immunogenic compositions  
15 and vaccines may be administered parenterally, by injection subcutaneously, intravenously, intradermally, intraperitoneally or intramuscularly, possibly following pretreatment of the injection site with a local anesthetic.

Alternatively, the immunogenic compositions formed according to the present invention, may be formulated and delivered in a manner to evoke an  
20 immune response at mucosal surfaces. Thus, the immunogenic composition may be administered to mucosal surfaces by, for example, the ocular, pulmonary, nasal or oral (intra-gastric) routes. Alternatively, other modes of administration including rectal, vaginal or urinary tract as well as suppositories may be desirable. For suppositories, binders and carriers may  
25 include, for example, polyalkylene glycols or triglycerides. Oral formulations may include normally employed excipients, such as, for example, pharmaceutical grades of saccharine, cellulose and magnesium carbonate.

The immunogenic preparations and vaccines are administered in a manner compatible with the dosage formulation, and in such amount as is  
30 therapeutically effective, protective and immunogenic. The quantity to be administered depends on the subject to be treated, including, for example, the capacity of the individual's immune system to synthesize the MOMP and

76 kDa proteins and antibodies thereto, and if needed, to produce a cell-mediated immune response. Precise amounts of active ingredient required to be administered depend on the judgement of the practitioner. However, suitable dosage ranges are readily determinable by one skilled in the art and  
5 may be of the order of about 1  $\mu$ g to about 1 mg of the vectors.

Suitable regimes for initial administration and booster doses are also variable, but may include an initial administration followed by subsequent administrations. The dosage may also depend on the route of administration and will vary according to the size of the host. A vaccine which protects  
10 against only one pathogen is a monovalent vaccine. Vaccines which contain antigenic material of several pathogens are combined vaccines and also belong to the present invention. Such combined vaccines contain, for example, material from various pathogens or from various strains of the same pathogen, or from combinations of various pathogens.

15 Immunogenicity may be significantly improved if the vectors are co-administered with adjuvants, commonly used as 0.05 to 0.1 percent solution in phosphate-buffered saline. Adjuvants enhance the immunogenicity of an antigen but are not necessarily immunogenic themselves. Adjuvants may act by retaining the antigen locally near the site of administration to produce  
20 a depot effect facilitating a slow, sustained release of antigen to cells of the immune system. Adjuvants can also attract cells of the immune system to an antigen depot and stimulate such cells to elicit immune responses.

Immunostimulatory agents or adjuvants have been used for many years to improve the host immune responses to, for example, vaccines.  
25 Thus, adjuvants have been identified that enhance the immune response to antigens. Some of these adjuvants are toxic, however, and can cause undesirable side-effects, making them unsuitable for use in humans and many animals. Indeed, only aluminum hydroxide and aluminum phosphate (collectively commonly referred to as alum) are routinely used as adjuvants  
30 in human and veterinary vaccines.

A wide range of extrinsic adjuvants and other immunomodulating material can provoke potent immune responses to antigens. These include

5 saponins complexed to membrane protein antigens to produce immune stimulating complexes (ISCOMS), pluronic polymers with mineral oil, killed mycobacteria in mineral oil, Freund's complete adjuvant, bacterial products, such as muramyl dipeptide (MDP) and lipopolysaccharide (LPS), as well as  
10 Quil A derivatives and components thereof, QS 21, calcium phosphate, calcium hydroxide, zinc hydroxide, an octodecyl ester of an amino acid, ISCOPREP, DC-chol, DDBA and polyphosphazene. Advantageous combinations of adjuvants are described in copending United States Patent Applications Nos.: 08/261,194 filed June 16, 1994 and 08/483,856 filed June  
15 7, 1995, assigned to the assignee hereof and the disclosures of which are incorporated herein by reference thereto (WO 95/34308).

In particular embodiments of the present invention, the vectors may be delivered in conjunction with a targeting molecule to target the vectors to selected cells including cells of the immune system.

15 The vectors may be delivered to the host by a variety of procedures, for example, Tang et al. (ref. 39) disclosed that introduction of gold microprojectiles coated with DNA encoding bovine growth hormone (BGH) into the skin of mice resulted in production of anti-BGH antibodies in the mice, while Furth et al. (ref. 40) showed that a jet injector could be used to  
20 transfect skin, muscle, fat and mammary tissues of living animals. See also U.S. Patents Nos. 4,245,050 and 5,015,580 and WO 94/24263.

#### **EXAMPLES**

The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the  
25 following specific examples. These examples are described solely for purposes of illustration and are not intended to limit the scope of the invention. Changes in form and substitution of equivalents are contemplated as circumstances may suggest or render expedient. Although specific terms have been employed herein, such terms are  
30 intended in a descriptive sense and not for purposes of limitation.



Example 1:

This Example illustrates the preparation of a plasmid vector pCA76kDa containing the 76kDa protein gene.

The 76kDa protein gene was amplified from *Chlamydia pneumoniae* (CM1) genomic DNA by polymerase chain reaction (PCR) using a 5' primer (5' GCTCTAGACCGCCATGACAAAAAACAT TATGCTTGGG 3') (SEQ ID No: 9) and 3' primer (5' CGGGATCCATAGAACTTGCTGCAGCGGG 3') (SEQ ID No: 10). The 5' primer contains a Xba I restriction site, a ribosome binding site, an initiation codon and a sequence close to the 5' end of the 76kDa protein coding sequence. The 3' primer includes the sequence encoding the C-terminal sequence of the 76kDa protein and a Bam HI restriction site. The stop codon was excluded and an additional nucleotide was inserted to obtain an inframe C-terminal fusion with the Histidine tag. The presence of a stop codon at nucleotide 828 of the amplified sequence means that only a partial 76kDa protein is expressed.

After amplification, the PCR fragment was using QIAquick™ PRC purification kit (Qiagen) and then digested with Xba I and Bam HI and cloned into the pCA-Myc-His eukaryotic expression vector as described in Example 3 below (Figure 2) with transcription under control of the human CMV promoter.

Example 2:

This Example illustrates the preparation of a plasmid vector pCAMOMP containing the MOMP protein gene.

The MOMP protein gene was amplified from *Chlamydia pneumoniae* (CM1) genomic DNA by polymerase chain reaction (PCR) using a 5' primer (5' CCCGGATATCCACCATGTTGCCTGTAGG GAACCCTTC 3') (SEQ ID No: 11) and a 3' primer (5' GGGGTACCGGAATCTGAACTGACCAGATACG 3') (SEQ ID No: 12). The 5' primer contains a EcoRV restriction site, a ribosome binding site, an initiation codon and a sequence encoding the N-terminal sequence of the mature MOMP. The 3' primer includes the sequence encoding the C-

terminal sequence of the MOMP and a Kpn I restriction site. The DNA sequence encoding the leader peptide was excluded, the stop codon was excluded and an additional nucleotide was inserted to obtain an in-frame C-terminal fusion with the Histidine tag.

5        After amplification, the PCR fragment was purified using QIAquick™ PCR purification kit (Qiagen) and then digested with Eco RV and Kpn I and cloned into the pCA-Myc-His eukaryotic expression vector described in Example 3 (Figure 4) with transcription under control of the human CMV promoter.

10    Example 3:

This Example illustrates the preparation of the eukaryotic expression vectors pCA76kDa and pCAMOMP.

15        Plasmid pcDNA3.1 (–) (Invitrogen) was restricted with Spe I and Bam HI to remove the CMV promoter and the remaining vector fragment was isolated. The CMV promoter and intron A from plasmid VR-1012 (Vical) was isolated on a Spe I / Bam HI fragment. The fragments were ligated together to produce plasmid pCA/Myc-His, as seen in Figure 2.

20        The Xba I/Bam HI restricted PCR fragment containing the 76kDa protein gene (Example 1) was ligated into the Xba I and Bam HI restricted plasmid pCA/Myc-His to produce plasmid pCA76kDa (Figure 2).

The Eco RV/Kpn I restricted PCR fragment containing the MOMP gene (Example 2) was ligated into Eco RV/Kpn I restricted pCA/Myc-His to produce plasmid pCAMOMP (Figure 4).

25        The resulting plasmids, pCA76kDa and pCAMOMP, were transferred by electroporation into *E. coli* XL-1 blue (Stratagene) which was grown in LB broth containing 50 µg/ml of carbenicillin. The plasmids were isolated by Endo Free Plasmid Giga Kit™ (Qiagen) large scale DNA purification system. DNA concentration was determined by absorbance at 260 nm and the plasmid was verified after gel electrophoresis and  
30    Ethidium bromide staining and comparison to molecular weight standards. The 5' and 3' ends of the gene were verified by sequencing using a LiCor model 4000 L DNA sequencer and IRD-800 labelled primers.

Example 4:

This Example illustrates the immunization of mice to achieve protection against an intranasal challenge by *C. pneumoniae*.

It has been previously demonstrated that mice are susceptible to intranasal infection with different isolates of *C. pneumoniae* (ref. 41). Strain AR-39 (ref. 42) was used in Balb/c mice as a challenge infection model to examine the capacity of chlamydia gene products delivered as naked DNA to elicit a protective response against a sublethal *C. pneumoniae* lung infection. Protective immunity is defined as an accelerated clearance of pulmonary infection.

Groups of 7 to 9 week old male Balb/c mice (5 to 9 per group) were immunized intramuscularly (i.m.) and intranasally (i.n.) with plasmids pCA76kDa and pCAMOMP containing the coding sequences of *C. pneumoniae* 76kDa and MOMP, respectively, prepared as described in Example 3. Saline or plasmid vectors containing non-protective inserted chlamydial genes, namely pCAI116 and pCAI178, were given to groups of control animals.

The constructs pCAI116 and pCAI178 are identical to pCA76kDa and pCAMOMP except that the nucleotide sequence encoding the partial 76kDa protein or MOMP is replaced with a *C. pneumoniae* nucleotide sequence encoding, respectively, a possible inclusion membrane protein and a nucleoside 5'-diphosphate phosphotransferase, respectively.

For i.m. immunization, alternate left and right quadriceps were injected with 100 µg of each DNA construct in 50 µl of PBS on three occasions at 0, 3, and 6 weeks. For i.n. immunization, anaesthetized mice aspirated 50 µl of PBS containing 50 µg of each DNA construct on three occasions at 0, 3, and 6 weeks. At week 8, immunized mice were inoculated i.n. with  $5 \times 10^5$  IFU of *C. pneumoniae*, strain AR39, in 100 µl of SPG buffer to test their ability to limit the growth of a sublethal *C. pneumoniae* challenge.

Lungs were taken from mice at day 9 post-challenge and immediately homogenized in SPG buffer (7.5% sucrose, 5 mM glutamate,

12.5 mM phosphate, pH 7.5). The homogenate was stored frozen at –70°C until assay. Dilutions of the homogenate were assayed for the presence of infectious chlamydia by inoculation onto monolayers of susceptible cells. The inoculum was centrifuged onto the cells at 3000 rpm for 1 hour, then the cells were incubated for three days at 35°C in the presence of 1 µg/ml cycloheximide. After incubation, the monolayers were fixed with formalin and methanol, then immunoperoxidase stained for the presence of *Chlamydial* inclusions using convalescent sera from rabbits infected with *C. pneumoniae* and metal-enhanced DAB as a peroxidase substrate.

Figure 5 and Table 1 contain the results obtained and show that mice immunized i.n. and i.m. with both pCA76kDa and pCAMOMP had chlamydial lung titers less than 6700 in 6 of 6 cases, whereas the range of values for control mice with saline were 15,000 to 106,100 IFU/lung in 20 out of 23 cases (mean 49,000) and 12,600 to 80,600 IFU/lung in 11 out of 12 cases (mean 33,500 to 47,000) for mice immunized with the vectors containing non-protective genes (Table 1). The mice immunized with only the pCAMOMP alone showed lung titres in the range of 5800 to 18,700 in 5 out of 6 cases (mean 12,600) and mice immunized with pCA76kDa alone showed similar titres in the range of 6,300 to 18,200 in 5 out of 6 cases (mean 7,400). The increased protection afforded by the combination of the two constructs is surprising in light of other failures due to antigen competition.

Table 1

MOUSE	BACTERIAL LOAD (INCLUSION FORMING UNITS PER LUNG) IN THE LUNGS OF BALB/C MICE IMMUNIZED WITH VARIOUS DNA IMMUNIZATION CONSTRUCTS					
	IMMUNIZING CONSTRUCT					
	Saline	pCAI116	pCAI178	pCAMOMP	pCA76kDa	pCAMOMP + pCA76kDa
	Day 9	Day 9	Day 9	Day 9	Day 9	Day 9
1	1700	47700	80600	5800	18200	6600
2	36200	12600	31900	30200	6300	5300
3	106100	28600	30600	9900	13400	0
4	33500	17700	6500	18700	100	3300
5	70400	77300	53000	0	2400	5200
6	48700	17600	79500	11000	4000	2700
7	600					
8	19800					
9	29500					
10	100000					
11	15000					
12	56600					
13	60300					
14	88800					
15	30400					
16	69300					
17	47500					
18	96500					
19	30200					
20	84800					
21	3800					
22	65900					
23	33000					
MEAN	49069.57	33583.33	47016.67	12600	7400	3850
SD	32120.48	24832.67	29524.32	10600.19	6981.40	2363.68

00916560 65/050" 90916560

SUMMARY OF THE DISCLOSURE

In summary of this disclosure, the present invention provides a novel immunization procedure for obtaining an enhanced protective immune response to Chlamydial infection by employing DNA immunization using nucleotide sequences encoding a MOMP and a 76 kDa protein of a strain of *Chlamydia*. Modifications are possible within the scope of the invention.

09391506 090799

# REFERENCES

1. Grayston et al. (1995), Journal of Infectious Diseases 168:1231
2. Campos et al. (1995), Investigation of Ophthalmology and Visual Science 36:1477
3. Grayston et al (1990), Journal of Infectious Diseases 161:618
4. Marrie (1993), Clinical Infectious Diseases. 18:501
5. Wang et al (1986), Chlamydial infections. Cambridge University Press, Cambridge. p. 329
6. Norman et al (1998), acta Paediatrica, 87:23-27
7. Saikku et al.(1988), Lancet;ii:983
8. Thom et al. (1992), JAMA 268:68
9. Linnanmaki et al. (1993), Circulation 87:1030
10. Saikku et al. (1992), Annals Internal Medicine 116:273
11. Melnick et al(1993), American Journal of Medicine 95:499
12. Shor et al. (1992), South African. Medical Journal 82:158
13. Kuo et al. (1993), Journal of Infectious Diseases 167:841
14. Kuo et al. (1993), Arteriosclerosis and Thrombosis 13:1500
15. Campbell et al (1995), Journal of Infectious Diseases 172:585
16. Chiu et al. Circulation, 1997 (In Press).
17. Ramirez et al (1996) Annals of Internal Medicine 125:979
18. Jackson et al. Abst. K121, p272, 36th ICAAC, 15-18 Sept. 1996, New Orleans.
19. Fong et al (1997) Journal of Clinical Microbiology 35:48
20. Hahn DL, et al. Evidence for Chlamydia pneumoniae infection in steroid-dependent asthma. Ann Allergy Asthma Immunol. 1998 Jan; 80(1): 45-49.

21. Hahn DL, et al. Association of Chlamydia pneumoniae IgA antibodies with recently symptomatic asthma. Epidemiol Infect. 1996 Dec; 117(3): 513-517.
22. Bjornsson E, et al. Serology of chlamydia in relation to asthma and bronchial hyperresponsiveness. Scand J Infect Dis. 1996; 28(1): 63-69.
23. Hahn DL. Treatment of *Chlamydia pneumoniae* infection in adult asthma: a before-after trial. J Fam Pract. 1995 Oct; 41(4): 345-351.
24. Allegra L, et al. Acute exacerbations of asthma in adults: role of Chlamydia pneumoniae infection. Eur Respir J. 1994 Dec; 7(12): 2165-2168.
25. Hahn DL, et al. Association of Chlamydia pneumoniae (strain TWAR) infection with wheezing, asthmatic bronchitis, and adult-onset asthma. JAMA. 1991 Jul 10; 266(2): 225-230.
26. Pal et al.(1996) Infection and Immunity.64:5341
27. Jones et al. (1995) Vaccine 13:715
28. Igietsemes et al. (1993) Immunology 5:317
29. Igietseme et al (1993) Regional Immunology 5:317
30. Magee et al (1993) Regional Immunology 5: 305
31. Landers et al (1991) Infection & Immunity 59:3774
32. Magee et al (1995) Infection & Immunity 63:516
33. Cotter et al. (1995) Infection and Immunity63:4704
34. Campbell et al (1990) Infection and Immunity 58:93
35. McCafferty et al (1995) Infection and Immunity 63:2387-9.
36. Knudsen et al (1996)Third Meeting of the European Society for Chlamydia Research, Vienna
37. Wiedmann-Al-Ahmad M, et al. Reactions of polyclonal and neutralizing anti-p54 monoclonal antibodies with an isolated, species-specific 54-kilodalton protein of Chlamydia pneumoniae. Clin Diagn Lab Immunol. 1997 Nov; 4(6): 700-704
38. Xiang, 1995, Immunity 2: 129-35.



39. Tang et al, Nature 1992; 356:152-154.
40. Furth et al., Vaccine 1994, 12:1503-1509.
41. Yang et al, 1993, Infection & Immunity, vol. 61, pp 2037-40.
42. Chi E.Y., Kuo C.C., Grayston J.T. 1987. Unique ultrastructure in the elementary body of Chlamydia sp strain TWAR. J. Bacteriol 169(8): 3757-63.

09391606 1090799

CLAIMS

What we claim is:

1. An immunogenic composition for *in vivo* administration to a host for the generation in the host of a protective immune response against Chlamydial infection, comprising:

a first vector comprising:

a first nucleotide sequence encoding a major outer membrane protein (MOMP) of a strain of *Chlamydia* and  
a first promoter sequence operatively coupled to said first nucleotide sequence for expression of said MOMP in the host;

a second vector comprising:

a second nucleotide sequence encoding a 76 kDa protein of a strain of *Chlamydia* and a second promoter sequence operatively coupled to said second nucleotide sequence for expression of said 76 kDa protein in the host; and

a pharmaceutically-acceptable carrier therefor.

2. The immunogenic composition of claim 1 wherein the first nucleotide sequence encodes a MOMP from *Chlamydia pneumoniae*.

3. The immunogenic composition of claim 1 wherein the first nucleotide sequence encodes a MOMP from *Chlamydia trachomatis*.

4. The immunogenic composition of claim 2 wherein said first nucleotide sequence has SEQ ID No: 12, 13 or 14.

5. The immunogenic composition of claim 2 wherein said first nucleotide sequence encodes a MOMP having SEQ ID No: 15 or 16.

6. The immunogenic composition of claim 2 wherein the first promoter is a cytomegalovirus promoter.

7. The immunogenic composition of claim 1 wherein the second nucleotide sequence encodes a 76 kDa protein from *Chlamydia pneumoniae*.

8. The immunogenic composition of claim 1 wherein the second nucleotide sequence encodes a 76 kDa protein from *Chlamydia trachomatis*.
9. The immunogenic composition of claim 7 wherein said second nucleotide sequence has SEQ ID No: 1, 2, 3 or 4.
10. The immunogenic composition of claim 7 wherein said second nucleotide sequence encodes a 76 kDa protein having a molecular size of about 35 kDa and having SEQ ID No: 7.
11. The immunogenic composition of claim 7 wherein said second nucleotide sequence encodes a 76 kDa protein having a molecular size of about 60 kDa and having SEQ ID No: 8 or 9.
12. The immunogenic composition of claim 7 wherein said second promoter is a cytomegalovirus promoter.
13. The immunogenic composition of claim 1 wherein said first vector is a plasmid vector.
14. The immunogenic composition of claim 13 wherein said first plasmid vector has the identifying characteristics of pCAMOMP as seen in Figure 4.
15. The immunogenic composition of claim 1 wherein said second vector is a plasmid vector.
16. The immunogenic composition of claim 15 wherein said second plasmid vector has the identifying characteristics of pCA76kDa as seen in Figure 2.
17. The immunogenic composition of claim 1 wherein both said first and second vectors are plasmid vectors.
18. The immunogenic composition of claim 17 wherein said first plasmid vector is pCAMOMP and said second plasmid vector is pCA76kDa.
19. The immunogenic composition of claim 1 wherein said first and second vectors are present in amounts such that the individual protective effect of each vector upon administration of the composition to the host is not adversely affected by the other.

20. The immunogenic composition of claim 1 wherein said first and second vectors are present in amounts such that an enhanced protective effect is achieved in comparison to the individual vectors alone.
21. A method of immunizing a host against disease caused by infection with a strain of *Chlamydia*, which comprises administering to said host an effective amount of an immunogenic composition of claim 1.✓
22. The method of claim 21 wherein said immunogenic composition is administered intranasally.
23. The method of claim 21 wherein said host is a human host.

ABSTRACT OF THE DISCLOSURE

A protective immune response against Chlamydial infection is achieved by *in vivo* administration of an immunogenic composition comprising two vectors and a pharmaceutically-acceptable carrier therefor.

- 5 One of the vectors comprises a first nucleotide sequence encoding a major outer membrane protein (MOMP) of a strain of *Chlamydia*, preferably *C. pneumoniae*, and a promoter sequence operatively coupled to the first nucleotide sequence for expression of the MOMP in the host. The other of the vectors comprises a second nucleotide sequence
- 10 encoding a 76 kDa protein of a strain of *Chlamydia*, preferably *C. pneumoniae*, and a promoter sequence operatively coupled to the second nucleotide sequence for expression of the 76 kDa protein in the host. The protection efficiency which is achieved by the immunization procedure is enhanced over that attained with the individual vectors alone.

Figure 1 Nucleotide Sequence of the 76kDa *C. pneumoniae* gene

CDS 5'

(175) .. (825)

CDS 3'

(940) .. (2409)

```

ttgcggtgct gttaacggtg gagggcagtg tagtctgagc agtactcggt gctgccgcgc 60
gcgccaccag acataatagc tgacagacta acagactggt cctttccatg ggtcttttct 120
gcagtcaccg tcgtcgacac gtgtgatcag atatcgcggc cgctctagac cgcc atg 177
                                     Met
                                     1

aca aaa aaa cat tat gct tgg gtt gta gaa ggg att ctc aat cgt ttg 225
Thr Lys Lys His Tyr Ala Trp Val Val Glu Gly Ile Leu Asn Arg Leu
          5                      10                      15

cct aaa cag ttt ttt gtg aaa tgt agt gtt gtc gac tgg aac aca ttc 273
Pro Lys Gln Phe Phe Val Lys Cys Ser Val Val Asp Trp Asn Thr Phe
          20                      25                      30

gtt cct tca gaa acc tcc act aca gaa aaa gct gct aca aac gct atg 321
Val Pro Ser Glu Thr Ser Thr Thr Glu Lys Ala Ala Thr Asn Ala Met
          35                      40                      45

aaa tac aaa tac tgt gtt tgg cag tgg ctc gtc gga aag cat agt cag 369
Lys Tyr Lys Tyr Cys Val Trp Gln Trp Leu Val Gly Lys His Ser Gln
          50                      55                      60                      65

gtt cct tgg atc aat gga cag aaa aag cct cta tat ctt tat gga gct 417
Val Pro Trp Ile Asn Gly Gln Lys Lys Pro Leu Tyr Leu Tyr Gly Ala
          70                      75                      80

ttc tta atg aac cct tta gca aag gct acg aag act acg tta aat gga 465
Phe Leu Met Asn Pro Leu Ala Lys Ala Thr Lys Thr Thr Leu Asn Gly
          85                      90                      95

aaa gaa aac cta gct tgg ttt att gga gga act tta ggg gga ctc aga 513
Lys Glu Asn Leu Ala Trp Phe Ile Gly Gly Thr Leu Gly Gly Leu Arg
          100                      105                      110

aaa gct gga gac tgg tct gcc aca gta cgt tat gag tat gtc gaa gcc 561
Lys Ala Gly Asp Trp Ser Ala Thr Val Arg Tyr Glu Tyr Val Glu Ala
          115                      120                      125

ttg tca gtt cca gaa ata gat gtt tca ggg att ggc cgt ggt aat tta 609
Leu Ser Val Pro Glu Ile Asp Val Ser Gly Ile Gly Arg Gly Asn Leu
          130                      135                      140                      145

tta aag ttt tgg ttc gcc caa gca att gct gct aac tat gat cct aaa 657
Leu Lys Phe Trp Phe Ala Gln Ala Ile Ala Ala Asn Tyr Asp Pro Lys

```


150	155	160	
gag gct aat agt ttt aca aat tat aaa gga ttt tcc gct cta tat atg			705
Glu Ala Asn Ser Phe Thr Asn Tyr Lys Gly Phe Ser Ala Leu Tyr Met			
165	170	175	
tat ggc atc aca gat tct cta tca ttc aga gct tat ggg gct tac tcc			753
Tyr Gly Ile Thr Asp Ser Leu Ser Phe Arg Ala Tyr Gly Ala Tyr Ser			
180	185	190	
aaa cca gca aac gat aaa ctc ggc agt gat ttt act ttc cga aag ttt			801
Lys Pro Ala Asn Asp Lys Leu Gly Ser Asp Phe Thr Phe Arg Lys Phe			
195	200	205	
gat cta ggt ata att tca gcg ttt taagtcaaattttaataaaaa tcttttaaaaa			855
Asp Leu Gly Ile Ile Ser Ala Phe			
210	215		
caggctcgca ttaattatta gtgagagctt tttttttatt ttttataata aaactaaaag			915
attttttatta ttttttgagt tttt atg gtt aat cct att ggt cca ggt cct			966
Met Val Asn Pro Ile Gly Pro Gly Pro			
220	225		
ata gac gaa aca gaa cgc aca cct ccc gca gat ctt tct gct caa gga			1014
Ile Asp Glu Thr Glu Arg Thr Pro Pro Ala Asp Leu Ser Ala Gln Gly			
230	235	240	
ttg gag gcg agt gca gca aat aag agt gcg gaa gct caa aga ata gca			1062
Leu Glu Ala Ser Ala Ala Asn Lys Ser Ala Glu Ala Gln Arg Ile Ala			
245	250	255	
ggt gcg gaa gct aag cct aaa gaa tct aag acc gat tct gta gag cga			1110
Gly Ala Glu Ala Lys Pro Lys Glu Ser Lys Thr Asp Ser Val Glu Arg			
260	265	270	
tgg agc atc ttg cgt tct gca gtg aat gct ctc atg agt ctg gca gat			1158
Trp Ser Ile Leu Arg Ser Ala Val Asn Ala Leu Met Ser Leu Ala Asp			
275	280	285	290
aag ctg ggt att gct tct agt aac agc tcg tct tct act agc aga tct			1206
Lys Leu Gly Ile Ala Ser Ser Asn Ser Ser Ser Ser Thr Ser Arg Ser			
295	300	305	
gca gac gtg gac tca acg aca gcg acc gca cct acg cct cct cca ccc			1254
Ala Asp Val Asp Ser Thr Thr Ala Thr Ala Pro Thr Pro Pro Pro Pro			
310	315	320	
acg tct gat gat tat aag act caa gcg caa aca gct tac gat act atc			1302
Thr Ser Asp Asp Tyr Lys Thr Gln Ala Gln Thr Ala Tyr Asp Thr Ile			
325	330	335	
ttt acc tca aca tca cta gct gac ata cag gct gct ttg gtg agc ctc			1350
Phe Thr Ser Thr Ser Leu Ala Asp Ile Gln Ala Ala Leu Val Ser Leu			

340	345	350	
cag gat gct gtc act aat ata aag gat aca gcg gct act gat gag gaa			1398
Gln Asp Ala Val Thr Asn Ile Lys Asp Thr Ala Ala Thr Asp Glu Glu			
355	360	365	370
acc gca atc gct gcg gag tgg gaa act aag aat gcc gat gca att aaa			1446
Thr Ala Ile Ala Ala Glu Trp Glu Thr Lys Asn Ala Asp Ala Ile Lys			
	375	380	385
gtt ggc gcg caa att aca gaa tta gcg aaa tat gct tcg gat aac caa			1494
Val Gly Ala Gln Ile Thr Glu Leu Ala Lys Tyr Ala Ser Asp Asn Gln			
	390	395	400
gcg att ctt gac tct tta ggt aaa ctg act tcc ttc gac ctc tta cag			1542
Ala Ile Leu Asp Ser Leu Gly Lys Leu Thr Ser Phe Asp Leu Leu Gln			
	405	410	415
act gct ctt ctc caa tct gta gca aac aat aac aaa gca gct gag ctt			1590
Thr Ala Leu Leu Gln Ser Val Ala Asn Asn Asn Lys Ala Ala Glu Leu			
	420	425	430
ctt aaa gag atg caa gat aac cca gta gtc cca ggg aaa acg cct gca			1638
Leu Lys Glu Met Gln Asp Asn Pro Val Val Pro Gly Lys Thr Pro Ala			
	435	440	445
att gct caa tct tta gtt gat cag aca gat gct aca gcg aca cag ata			1686
Ile Ala Gln Ser Leu Val Asp Gln Thr Asp Ala Thr Ala Thr Gln Ile			
	455	460	465
gag aaa gat gga aat gcg att ggg gat gca tat ttt gca gga cag aac			1734
Glu Lys Asp Gly Asn Ala Ile Gly Asp Ala Tyr Phe Ala Gly Gln Asn			
	470	475	480
gct agt gga gct gta gaa aat gct aaa tct aat aac agt ata agc aac			1782
Ala Ser Gly Ala Val Glu Asn Ala Lys Ser Asn Asn Ser Ile Ser Asn			
	485	490	495
ata gat tca gct aaa gca gca atc gct act gct aag aca caa ata gct			1830
Ile Asp Ser Ala Lys Ala Ala Ile Ala Thr Ala Lys Thr Gln Ile Ala			
	500	505	510
gaa gct cag aaa aag ttc ccc gac tct cca att ctt caa gaa gcg gaa			1878
Glu Ala Gln Lys Lys Phe Pro Asp Ser Pro Ile Leu Gln Glu Ala Glu			
	515	520	525
caa atg gta ata cag gct gag aaa gat ctt aaa aat atc aaa cct gca			1926
Gln Met Val Ile Gln Ala Glu Lys Asp Leu Lys Asn Ile Lys Pro Ala			
	535	540	545
gat ggt tct gat gtt cca aat cca gga act aca gtt gga ggc tcc aag			1974
Asp Gly Ser Asp Val Pro Asn Pro Gly Thr Thr Val Gly Gly Ser Lys			
	550	555	560



caa caa gga agt agt att ggt agt att cgt gtt tcc atg ctg tta gat	2022
Gln Gln Gly Ser Ser Ile Gly Ser Ile Arg Val Ser Met Leu Leu Asp	
565 570 575	
gat gct gaa aat gag acc gct tcc att ttg atg tct ggg ttt cgt cag	2070
Asp Ala Glu Asn Glu Thr Ala Ser Ile Leu Met Ser Gly Phe Arg Gln	
580 585 590	
atg att cac atg ttc aat acg gaa aat cct gat tct caa gct gcc caa	2118
Met Ile His Met Phe Asn Thr Glu Asn Pro Asp Ser Gln Ala Ala Gln	
595 600 605 610	
cag gag ctc gca gca caa gct aga gca gcg aaa gcc gct gga gat gac	2166
Gln Glu Leu Ala Ala Gln Ala Arg Ala Ala Lys Ala Ala Gly Asp Asp	
615 620 625	
agt gct gct gca gcg ctg gca gat gct cag aaa gct tta gaa gcg gct	2214
Ser Ala Ala Ala Ala Leu Ala Asp Ala Gln Lys Ala Leu Glu Ala Ala	
630 635 640	
cta ggt aaa gct ggg caa caa cag ggc ata ctc aat gct ttg gga cag	2262
Leu Gly Lys Ala Gly Gln Gln Gln Gly Ile Leu Asn Ala Leu Gly Gln	
645 650 655	
atc gct tct gct gct gtt gtg agc gca gga gtc ctc ccg ctg cag caa	2310
Ile Ala Ser Ala Ala Val Val Ser Ala Gly Val Leu Pro Leu Gln Gln	
660 665 670	
gtt cta tgg atc cga gct cgg tac caa gct tac gta gaa caa aaa ctc	2358
Val Leu Trp Ile Arg Ala Arg Tyr Gln Ala Tyr Val Glu Gln Lys Leu	
675 680 685 690	
atc tca gaa gag gat ctg aat agc gcc gtc gac cat cat cat cat cat	2406
Ile Ser Glu Glu Asp Leu Asn Ser Ala Val Asp His His His His His	
695 700 705	
cat tgagttttaa cggctctccag cttaagttta aaccgctgat cagcctcgac	2459
<u>His</u>	
tgtgccttct agttgccagc catctgttgt ttgccccctcc cccgtgcctt ccttgaccct	2519
ggaaggtgcc actcccactg tccttt	2545

Figure 2 Construction of pCAD76kDa



039166 090799 65/060" 909T6E20

Figure 3 Nucleotide sequence of the *C. pneumoniae* MOMP gene.

(126) .. (1307)

```

tgagcagtac tcgttgctgc cgcgcgcgcc accagacata atagctgaca gactaacaga 60
ctgttccttt ccatgggtct tttctgcagt caccgtcgtc gacacgtgtg atcagatatc 120
ccacc atg ttg cct gta ggg aac cct tct gat cca agc tta tta att gat 170
    Met Leu Pro Val Gly Asn Pro Ser Asp Pro Ser Leu Leu Ile Asp
        1             5             10             15

ggt aca ata tgg gaa ggt gct gca gga gat cct tgc gat cct tgc gct 218
Gly Thr Ile Trp Glu Gly Ala Ala Gly Asp Pro Cys Asp Pro Cys Ala
        20             25             30

act tgg tgc gac gct att agc tta cgt gct gga ttt tac gga gac tat 266
Thr Trp Cys Asp Ala Ile Ser Leu Arg Ala Gly Phe Tyr Gly Asp Tyr
        35             40             45

gtt ttc gac cgt atc tta aaa gta gat gca cct aaa aca ttt tct atg 314
Val Phe Asp Arg Ile Leu Lys Val Asp Ala Pro Lys Thr Phe Ser Met
        50             55             60

gga gcc aag cct act gga tcc gct gct gca aac tat act act gcc gta 362
Gly Ala Lys Pro Thr Gly Ser Ala Ala Ala Asn Tyr Thr Thr Ala Val
        65             70             75

gat aga cct aac ccg gcc tac aat aag cat tta cac gat gca gag tgg 410
Asp Arg Pro Asn Pro Ala Tyr Asn Lys His Leu His Asp Ala Glu Trp
        80             85             90             95

ttc act aat gca ggc ttc att gcc tta aac att tgg gat cgc ttt gat 458
Phe Thr Asn Ala Gly Phe Ile Ala Leu Asn Ile Trp Asp Arg Phe Asp
        100             105             110

gtt ttc tgt act tta gga gct tct aat ggt tac att aga gga aac tct 506
Val Phe Cys Thr Leu Gly Ala Ser Asn Gly Tyr Ile Arg Gly Asn Ser
        115             120             125

aca gcg ttc aat ctc gtt ggt tta ttc gga gtt aaa ggt act act gta 554
Thr Ala Phe Asn Leu Val Gly Leu Phe Gly Val Lys Gly Thr Thr Val
        130             135             140

aat gca aat gaa cta cca aac gtt tct tta agt aac gga gtt gtt gaa 602
Asn Ala Asn Glu Leu Pro Asn Val Ser Leu Ser Asn Gly Val Val Glu
        145             150             155

ctt tac aca gac acc tct ttc tct tgg agc gta ggc gct cgt gga gcc 650
Leu Tyr Thr Asp Thr Ser Phe Ser Trp Ser Val Gly Ala Arg Gly Ala
        160             165             170             175

```

00991606"090799

tta tgg gaa tgc ggt tgt gca act ttg gga gct gaa ttc caa tat gca	698
Leu Trp Glu Cys Gly Cys Ala Thr Leu Gly Ala Glu Phe Gln Tyr Ala	
180 185 190	
cag tcc aaa cct aaa gtt gaa gaa ctt aat gtg atc tgt aac gta tcg	746
Gln Ser Lys Pro Lys Val Glu Glu Leu Asn Val Ile Cys Asn Val Ser	
195 200 205	
caa ttc tct gta aac aaa ccc aag ggc tat aaa ggc gtt gct ttc ccc	794
Gln Phe Ser Val Asn Lys Pro Lys Gly Tyr Lys Gly Val Ala Phe Pro	
210 215 220	
ttg cca aca gac gct ggc gta gca aca gct act gga aca aag tct gcg	842
Leu Pro Thr Asp Ala Gly Val Ala Thr Ala Thr Gly Thr Lys Ser Ala	
225 230 235	
acc atc aat tat cat gaa tgg caa gta gga gcc tct cta tct tac aga	890
Thr Ile Asn Tyr His Glu Trp Gln Val Gly Ala Ser Leu Ser Tyr Arg	
240 245 250 255	
cta aac tct tta gtg cca tac att gga gta caa tgg tct cga gca act	938
Leu Asn Ser Leu Val Pro Tyr Ile Gly Val Gln Trp Ser Arg Ala Thr	
260 265 270	
ttt gat gct gat aac atc cgc att gct cag cca aaa cta cct aca gct	986
Phe Asp Ala Asp Asn Ile Arg Ile Ala Gln Pro Lys Leu Pro Thr Ala	
275 280 285	
gtt tta aac tta act gca tgg aac cct tct tta cta gga aat gcc aca	1034
Val Leu Asn Leu Thr Ala Trp Asn Pro Ser Leu Leu Gly Asn Ala Thr	
290 295 300	
gca ttg tct act act gat tcg ttc tca gac ttc atg caa att gtt tcc	1082
Ala Leu Ser Thr Thr Asp Ser Phe Ser Asp Phe Met Gln Ile Val Ser	
305 310 315	
tgt cag atc aac aag ttt aaa tct aga aaa gct tgt gga gtt act gta	1130
Cys Gln Ile Asn Lys Phe Lys Ser Arg Lys Ala Cys Gly Val Thr Val	
320 325 330 335	
gga gct act tta gtt gat gct gat aaa tgg tca ctt act gca gaa gct	1178
Gly Ala Thr Leu Val Asp Ala Asp Lys Trp Ser Leu Thr Ala Glu Ala	
340 345 350	
cgt tta att aac gag aga gct gct cac gta tct ggt cag ttc aga ttc	1226
Arg Leu Ile Asn Glu Arg Ala Ala His Val Ser Gly Gln Phe Arg Phe	
355 360 365	
cgg tac caa gct tac gta gaa caa aaa ctc atc tca gaa gag gat ctg	1274
Arg Tyr Gln Ala Tyr Val Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu	
370 375 380	
aat agc gcc gtc gac cat cat cat cat cat cat tgagtttaaa cggtctccag	1327
Asn Ser Ala Val Asp His His His His His His	

385


390

cttaagttta aaccgctgat cagcctcgac tgtgccttct agttgccagc catctgttgt 1387


ttgcccctcc cccgtgcctt ccttgaccct ggaaggtgc 1426

00391506.090799

Figure 4 Construction of pCAMOMP



093460" 909T6E60



Docket No.  
1038-971 MIS:jb

# Declaration and Power of Attorney For Patent Application

## English Language Declaration

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

**IMMUNOGENIC COMPOSITIONS FOR PROTECTION AGAINST CHLAMYDIAL INFECTION**

the specification of which

(check one)

☒ is attached hereto.

☐ was filed on \_\_\_\_\_ as United States Application No. or PCT International Application Number \_\_\_\_\_ and was amended on \_\_\_\_\_ (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Priority Not Claimed

_____ (Number)	_____ (Country)	_____ (Day/Month/Year Filed)	<input type="checkbox"/>
_____ (Number)	_____ (Country)	_____ (Day/Month/Year Filed)	<input type="checkbox"/>
_____ (Number)	_____ (Country)	_____ (Day/Month/Year Filed)	<input type="checkbox"/>



I hereby claim the benefit under 35 U.S.C. Section 119(e) of any United States provisional application(s) listed below:

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

I hereby claim the benefit under 35 U. S. C. Section 120 of any United States application(s), or Section 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. Section 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, C. F. R., Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application:

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Status)  
(patented, pending, abandoned)

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Status)  
(patented, pending, abandoned)

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Status)  
(patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. *(list name and registration number)*

**Michael I. Stewart (24,973)**

Send Correspondence to: **Sim & McBurney**  
**6th Floor, 330 University Avenue**  
**Toronto, Ontario**  
**Canada, M5G 1R7.**

Direct Telephone Calls to: *(name and telephone number)*  
**(416) 595-1155**

Full name of sole or first inventor

**Andrew D. Mordin**

Sole or first inventor's signature

Date

Residence

**Newmarket, Ontario, Canada**

Citizenship

**Canadian**

Post Office Address

**146 Rhodes Circle, Newmarket, Ontario, Canada, L3X 1V2.**

Full name of second inventor, if any

**Pamela L. Dunn**

Second inventor's signature

Date

Residence

**Mississauga, Ontario, Canada**

Citizenship

**Canadian**

Post Office Address

**Apt. 703, 3700 Kaneff Crescent, Mississauga, Ontario, Canada, L5A 4B8.**